

## 1-(5-Bromo-2-oxoindolin-3-ylidene)thiosemicarbazide acetonitrile monosolvate

Fernanda Rosi Soares Pederzolli,<sup>a</sup> Leandro Bresolin,<sup>a\*</sup>  
 Vanessa Santana Carratu,<sup>a</sup> Aline Locatelli<sup>b</sup> and  
 Adriano Bof de Oliveira<sup>c</sup>

<sup>a</sup>Escola de Química e Alimentos, Universidade Federal do Rio Grande, Av. Itália km 08, Campus Carreiros, 96201-900 Rio Grande, RS, Brazil, <sup>b</sup>Departamento de Química, Universidade Federal de Santa Maria, Av. Roraima, Campus, 97105-900 Santa Maria, RS, Brazil, and <sup>c</sup>Departamento de Química, Universidade Federal de Sergipe, Av. Marechal Rondon s/n, Campus, 49100-000 São Cristóvão-SE, Brazil  
 Correspondence e-mail: leandro\_bresolin@yahoo.com.br


Received 9 June 2011; accepted 17 June 2011

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.005\text{ \AA}$ ;  $R$  factor = 0.039;  $wR$  factor = 0.105; data-to-parameter ratio = 19.5.

In the crystal structure of the title compound,  $\text{C}_9\text{H}_7\text{BrN}_4\text{OS}\cdot\text{C}_2\text{H}_3\text{N}$ , the molecules are connected via  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{N}-\text{H}\cdots\text{S}$  interactions into zigzag chains perpendicular to [001]. The molecules in these chains are additionally linked to acetonitrile solvent molecules through  $\text{N}-\text{H}\cdots\text{N}$  hydrogen bonding. The molecules are arranged in layers and are stacked in the direction of the  $c$  axis indicative of  $\pi-\pi$  interactions, with distance = 3.381 (7) Å for the  $\text{C}\cdots\text{C}$  interaction parallel to [001]. An intramolecular  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bond is also observed in the main molecule.

### Related literature

For the pharmacological properties of isatin-thiosemicarbazone derivatives against cruzain, falcipain-2 and rhodesain, see: Chiyanzu *et al.* (2003). For the synthesis of 5-bromoisatin-3-thiosemicarbazone, see: Campaigne & Archer (1952).



### Experimental

#### Crystal data

$\text{C}_9\text{H}_7\text{BrN}_4\text{OS}\cdot\text{C}_2\text{H}_3\text{N}$	$V = 2729.6\text{ (12) \AA}^3$
$M_r = 340.21$	$Z = 8$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation
$a = 20.017\text{ (4) \AA}$	$\mu = 3.16\text{ mm}^{-1}$
$b = 13.352\text{ (2) \AA}$	$T = 293\text{ K}$
$c = 13.190\text{ (5) \AA}$	$0.22 \times 0.20 \times 0.16\text{ mm}$
$\beta = 129.258\text{ (2)\text{ }^\circ}$	

#### Data collection

Bruker CCD X8 APEXII diffractometer	3377 independent reflections
10884 measured reflections	2754 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.030$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	173 parameters
$wR(F^2) = 0.105$	H-atom parameters constrained
$S = 1.09$	$\Delta\rho_{\text{max}} = 1.03\text{ e \AA}^{-3}$
3377 reflections	$\Delta\rho_{\text{min}} = -0.86\text{ e \AA}^{-3}$

**Table 1**

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N3—H5···O	0.86	2.10	2.769 (3)	134
N4—H6···N5	0.86	2.61	3.438 (5)	161
N4—H7···O <sup>i</sup>	0.86	2.05	2.906 (4)	173
N1—H4···S <sup>ii</sup>	0.86	2.50	3.350 (3)	169

Symmetry codes: (i)  $x - \frac{1}{2}, -y - \frac{1}{2}, z - \frac{1}{2}$ ; (ii)  $x + \frac{1}{2}, -y - \frac{1}{2}, z + \frac{1}{2}$ .

Data collection: *APEX2* (Bruker, 2006); cell refinement: *SAINT* (Bruker, 2003); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We gratefully acknowledge Professor Dr Manfredo Hörner (Department of Chemistry, Federal University of Santa Maria, Brazil) for his help and support with the X-ray measurements, and CNPq/FAPERGS for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: NC2233).

### References

- Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2003). *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Bruker (2006). *APEX2*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Campaigne, E. & Archer, W. L. (1952). *J. Am. Chem. Soc.* **74**, 5801.
- Chiyanzu, I., Hansell, E., Gut, J., Rosenthal, P. J., McKerrow, J. H. & Chibale, K. (2003). *Bioorg. Med. Chem. Lett.* **13**, 3527–3530.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

## **supplementary materials**

*Acta Cryst.* (2011). E67, o1804 [doi:10.1107/S1600536811023786]

## 1-(5-Bromo-2-oxoindolin-3-ylidene)thiosemicarbazide acetonitrile monosolvate

F. R. S. Pederzolli, L. Bresolin, V. S. Carratu, A. Locatelli and A. B. de Oliveira

### Comment

Thiosemicarbazone derivatives have a wide range of biological properties. For example, isatin-based synthetic thiosemicarbazones show pharmacological activity against cruzain, falcipain-2 and rhodesain (Chiyanzu *et al.*, 2003). As part of our study of thiosemicarbazone derivatives, we report herein the crystal structure of 5-Bromoisatin-3-thiosemicarbazone acetonitrile solvate.

The crystal structure of the title compound is build of one-dimensional zigzag chain in which the molecules are linked by pairs of N—H···O and N—H···S hydrogen bonding. Each two molecules within these chains are additionally linked by acetonitrile molecules *via* N—H···N hydrogen bonding and weak C—H···S interactions. The molecules are arranged in layers and are stacked into the direction of the *c*-axis indicative for  $\pi$ – $\pi$ -interactions.

### Experimental

Starting materials were commercially available and were used without further purification. The synthesis was adapted from a procedure reported previously (Campaigne & Archer, 1952). The hydrochloric acid catalyzed reaction of 5-bromoisatin (8.83 mmol) and thiosemicarbazide (8.83 mmol) in ethanol (50 ml) was refluxed for 6 h. After cooling and filtering, crystals suitable for X-ray diffraction were obtained from an acetonitrile solution.

### Refinement

The C—H and N—H H atoms were positioned with idealized geometry and were refined isotropic with  $U_{\text{eq}}(\text{H})$  set to 1.2 times of the  $U_{\text{eq}}$  of the parent atom (1.5 for methyl H atoms) using a riding model with C—H = 0.93 Å for aromatic), C—H = 0.96 Å for methyl and N—H = 0.86 Å for N—H H atoms.

### Figures




Fig. 1. : The molecular structure of the title compound with labelling and displacement ellipsoids drawn at the 50% probability level.




Fig. 2. : Crystal structure of the title compound viewed in the direction of the crystallographic *c* axis. Hydrogen bonding is indicated as dashed lines.

# supplementary materials

---

## 1-(5-Bromo-2-oxoindolin-3-ylidene)thiosemicarbazide acetonitrile monosolvate

### Crystal data

C <sub>9</sub> H <sub>7</sub> BrN <sub>4</sub> OS·C <sub>2</sub> H <sub>3</sub> N	F(000) = 1360
M <sub>r</sub> = 340.21	D <sub>x</sub> = 1.656 Mg m <sup>-3</sup>
Monoclinic, C2/c	Melting point: 544.15 K
Hall symbol: -C 2yc	Mo K $\alpha$ radiation, $\lambda$ = 0.71073 Å
a = 20.017 (4) Å	Cell parameters from 3718 reflections
b = 13.352 (2) Å	$\theta$ = 2.6–27.6°
c = 13.190 (5) Å	$\mu$ = 3.16 mm <sup>-1</sup>
$\beta$ = 129.258 (2)°	T = 293 K
V = 2729.6 (12) Å <sup>3</sup>	Block, yellow
Z = 8	0.22 × 0.20 × 0.16 mm

### Data collection

Bruker CCD X8 APEXII diffractometer	2754 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube, Bruker CCD X8 APEXII	$R_{\text{int}} = 0.030$
graphite	$\theta_{\text{max}} = 28.3^\circ$ , $\theta_{\text{min}} = 2.0^\circ$
$\varphi$ and $\omega$ scans	$h = -26 \rightarrow 14$
10884 measured reflections	$k = -16 \rightarrow 17$
3377 independent reflections	$l = -7 \rightarrow 17$

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.039$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.105$	H-atom parameters constrained
$S = 1.09$	$w = 1/[\sigma^2(F_o^2) + (0.0396P)^2 + 12.0809P]$ where $P = (F_o^2 + 2F_c^2)/3$
3377 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
173 parameters	$\Delta\rho_{\text{max}} = 1.03 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.86 \text{ e \AA}^{-3}$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C9	0.3611 (2)	-0.2695 (2)	0.4897 (3)	0.0175 (6)
C10	0.0960 (2)	-0.0277 (3)	0.2196 (3)	0.0297 (8)
H8	0.0967	-0.0995	0.2159	0.045*
H9	0.0735	-0.0003	0.1361	0.045*
H10	0.0602	-0.0075	0.2410	0.045*
C11	0.1832 (2)	0.0088 (3)	0.3192 (4)	0.0296 (8)
C6	0.52345 (19)	0.0354 (2)	0.6478 (3)	0.0166 (6)
C5	0.4655 (2)	0.1144 (2)	0.5889 (3)	0.0169 (6)
H3	0.4062	0.1038	0.5301	0.020*
C4	0.5007 (2)	0.2109 (2)	0.6225 (3)	0.0205 (6)
C3	0.5890 (2)	0.2282 (2)	0.7097 (3)	0.0214 (7)
H2	0.6101	0.2934	0.7300	0.026*
C2	0.6469 (2)	0.1459 (3)	0.7675 (3)	0.0224 (7)
H1	0.7063	0.1558	0.8254	0.027*
C1	0.6126 (2)	0.0514 (2)	0.7353 (3)	0.0181 (6)
C8	0.5995 (2)	-0.1188 (3)	0.7243 (3)	0.0191 (6)
C7	0.51065 (19)	-0.0725 (2)	0.6367 (3)	0.0152 (6)
Br	0.42473 (2)	0.32205 (3)	0.54837 (4)	0.02739 (12)
N2	0.43754 (17)	-0.1178 (2)	0.5672 (2)	0.0171 (5)
N3	0.43806 (16)	-0.2191 (2)	0.5678 (2)	0.0174 (5)
H5	0.4860	-0.2515	0.6165	0.021*
N4	0.29129 (18)	-0.2149 (2)	0.4244 (3)	0.0217 (6)
H6	0.2951	-0.1507	0.4317	0.026*
H7	0.2414	-0.2430	0.3741	0.026*
N5	0.2515 (2)	0.0381 (3)	0.3981 (4)	0.0491 (10)
N1	0.65560 (17)	-0.0418 (2)	0.7786 (3)	0.0197 (5)
H4	0.7108	-0.0486	0.8329	0.024*
O	0.61642 (14)	-0.20772 (17)	0.7432 (2)	0.0200 (5)
S	0.36470 (5)	-0.39515 (6)	0.48661 (8)	0.02291 (19)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C9	0.0139 (15)	0.0224 (17)	0.0165 (14)	-0.0033 (12)	0.0098 (13)	-0.0014 (12)
C10	0.0273 (19)	0.0265 (19)	0.0264 (18)	-0.0018 (15)	0.0126 (16)	-0.0003 (15)
C11	0.0214 (18)	0.0231 (18)	0.035 (2)	0.0042 (14)	0.0134 (17)	0.0086 (15)
C6	0.0129 (14)	0.0219 (16)	0.0136 (14)	-0.0017 (12)	0.0077 (12)	-0.0009 (12)
C5	0.0157 (15)	0.0187 (15)	0.0156 (14)	0.0004 (12)	0.0097 (13)	0.0004 (12)
C4	0.0239 (17)	0.0175 (16)	0.0227 (16)	0.0048 (13)	0.0160 (15)	0.0041 (13)

## supplementary materials

---

C3	0.0248 (17)	0.0163 (16)	0.0259 (17)	-0.0049 (13)	0.0173 (15)	-0.0006 (13)
C2	0.0138 (15)	0.0238 (17)	0.0244 (17)	-0.0037 (13)	0.0096 (14)	-0.0032 (13)
C1	0.0149 (15)	0.0230 (16)	0.0164 (14)	-0.0006 (12)	0.0099 (13)	-0.0005 (12)
C8	0.0124 (15)	0.0271 (18)	0.0147 (14)	-0.0001 (13)	0.0072 (13)	0.0000 (12)
C7	0.0119 (14)	0.0188 (15)	0.0128 (13)	0.0001 (11)	0.0068 (12)	-0.0005 (11)
Br	0.0271 (2)	0.01948 (18)	0.0320 (2)	0.00440 (14)	0.01693 (16)	0.00453 (14)
N2	0.0164 (13)	0.0179 (13)	0.0172 (13)	0.0003 (10)	0.0107 (12)	0.0007 (10)
N3	0.0101 (12)	0.0179 (13)	0.0187 (13)	0.0011 (10)	0.0066 (11)	0.0003 (10)
N4	0.0130 (13)	0.0167 (13)	0.0283 (15)	-0.0002 (11)	0.0097 (12)	0.0017 (11)
N5	0.0247 (19)	0.046 (2)	0.052 (2)	-0.0026 (16)	0.0130 (18)	0.0101 (19)
N1	0.0110 (12)	0.0191 (14)	0.0223 (13)	-0.0005 (10)	0.0073 (11)	-0.0002 (11)
O	0.0133 (11)	0.0176 (11)	0.0212 (11)	0.0014 (9)	0.0072 (10)	0.0005 (9)
S	0.0140 (4)	0.0176 (4)	0.0273 (4)	0.0001 (3)	0.0084 (3)	0.0001 (3)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

C9—N4	1.305 (4)	C3—C2	1.419 (5)
C9—N3	1.370 (4)	C3—H2	0.9300
C9—S	1.681 (3)	C2—C1	1.369 (5)
C10—C11	1.451 (5)	C2—H1	0.9300
C10—H8	0.9600	C1—N1	1.412 (4)
C10—H9	0.9600	C8—O	1.217 (4)
C10—H10	0.9600	C8—O	1.217 (4)
C11—N5	1.142 (5)	C8—N1	1.346 (4)
C11—N5	1.142 (5)	C8—C7	1.510 (4)
C6—C5	1.386 (4)	C7—N2	1.285 (4)
C6—C1	1.398 (4)	N2—N3	1.352 (4)
C6—C7	1.455 (4)	N3—H5	0.8600
C5—C4	1.399 (5)	N4—H6	0.8600
C5—H3	0.9300	N4—H7	0.8600
C4—C3	1.389 (5)	N1—H4	0.8600
C4—Br	1.894 (3)		
N4—C9—N3	116.5 (3)	C1—C2—H1	121.0
N4—C9—S	125.9 (2)	C3—C2—H1	121.0
N3—C9—S	117.6 (2)	C2—C1—C6	121.6 (3)
C11—C10—H8	109.5	C2—C1—N1	129.0 (3)
C11—C10—H9	109.5	C6—C1—N1	109.4 (3)
H8—C10—H9	109.5	O—C8—O	0.00 (5)
C11—C10—H10	109.5	O—C8—N1	127.3 (3)
H8—C10—H10	109.5	O—C8—C7	127.3 (3)
H9—C10—H10	109.5	O—C8—C7	126.6 (3)
N5—C11—N5	0.0 (8)	O—C8—C7	126.6 (3)
N5—C11—C10	179.2 (5)	N1—C8—C7	106.1 (3)
N5—C11—C10	179.2 (5)	N2—C7—C6	125.9 (3)
C5—C6—C1	121.7 (3)	N2—C7—C8	127.7 (3)
C5—C6—C7	131.7 (3)	C6—C7—C8	106.3 (3)
C1—C6—C7	106.6 (3)	C7—N2—N3	117.8 (3)
C6—C5—C4	116.6 (3)	N2—N3—C9	119.0 (3)
C6—C5—H3	121.7	N2—N3—H5	120.5

C4—C5—H3	121.7	C9—N3—H5	120.5
C3—C4—C5	122.5 (3)	C9—N4—H6	120.0
C3—C4—Br	118.8 (3)	C9—N4—H7	120.0
C5—C4—Br	118.6 (2)	H6—N4—H7	120.0
C4—C3—C2	119.6 (3)	C8—N1—C1	111.6 (3)
C4—C3—H2	120.2	C8—N1—H4	124.2
C2—C3—H2	120.2	C1—N1—H4	124.2
C1—C2—C3	117.9 (3)		
C1—C6—C5—C4	−0.6 (4)	O—C8—C7—N2	0.2 (5)
C7—C6—C5—C4	−180.0 (3)	N1—C8—C7—N2	−178.7 (3)
C6—C5—C4—C3	0.3 (4)	O—C8—C7—C6	179.1 (3)
C6—C5—C4—Br	−178.6 (2)	O—C8—C7—C6	179.1 (3)
C5—C4—C3—C2	0.3 (5)	N1—C8—C7—C6	0.2 (3)
Br—C4—C3—C2	179.2 (2)	C6—C7—N2—N3	179.4 (3)
C4—C3—C2—C1	−0.7 (5)	C8—C7—N2—N3	−1.9 (4)
C3—C2—C1—C6	0.4 (5)	C7—N2—N3—C9	−176.9 (3)
C3—C2—C1—N1	179.9 (3)	N4—C9—N3—N2	−3.6 (4)
C5—C6—C1—C2	0.3 (5)	S—C9—N3—N2	176.5 (2)
C7—C6—C1—C2	179.8 (3)	C10—C11—N5—N5	0(12)
C5—C6—C1—N1	−179.3 (3)	O—C8—N1—C1	−179.0 (3)
C7—C6—C1—N1	0.2 (3)	O—C8—N1—C1	−179.0 (3)
C5—C6—C7—N2	−1.8 (5)	C7—C8—N1—C1	−0.1 (3)
C1—C6—C7—N2	178.7 (3)	C2—C1—N1—C8	−179.6 (3)
C5—C6—C7—C8	179.2 (3)	C6—C1—N1—C8	0.0 (3)
C1—C6—C7—C8	−0.2 (3)	N1—C8—O—O	0.0 (2)
O—C8—C7—N2	0.2 (5)	C7—C8—O—O	0.00 (11)

*Hydrogen-bond geometry (Å, °)*

D—H···A	D—H	H···A	D···A	D—H···A
N3—H5···O	0.86	2.10	2.769 (3)	134
N4—H6···N5	0.86	2.61	3.438 (5)	161
N4—H7···O <sup>i</sup>	0.86	2.05	2.906 (4)	173
N1—H4···S <sup>ii</sup>	0.86	2.50	3.350 (3)	169

Symmetry codes: (i)  $x-1/2, -y-1/2, z-1/2$ ; (ii)  $x+1/2, -y-1/2, z+1/2$ .

## supplementary materials

---

Fig. 1




Fig. 2

